


ChloroSan

Cleaning and Disinfection

How to clean **Blood Spills**



Disposable gloves and apron are required at all times. If there is a risk of splashing, please use safety glasses.



Fill the dilution bottle with 500ml of **cold water**. Add 5 ChloroSan tablets (10,000ppm solution). The lid must be off until the tablets have completely dissolved and must then be replaced.




Ensure the spill area is secure. Before applying ChloroSan, place paper towels over the blood spill.



Pour sufficient ChloroSan 10,000ppm solution on to the spill to cover it, and leave for 3 minutes before disposing of the paper towels as clinical waste.



Clean spill area and immediate surroundings with a 10,000ppm ChloroSan solution (5 tablets per 500ml of water).



ChloroSan solution should be kept for max. 24 hours and any residual solution disposed of with cold running water. Rinse bottle. Dispose of PPE into correct waste stream and wash hands.

Product Name	Cleaning and Disinfection	Disinfection Only	Product Use	Tablet Use - 10,000ppm Solution
ChloroSan 1.7g+	Yes	No	Blood Spills Disinfection and Area Clean Up (10,000ppm) (Contact time 3 mins)	5 tablets per 500ml of water

When used to manufacturer's instructions, this product complies with EN13727, EN13624, EN13697, EN14476, EN16777, and EN17126. A Technical Data and Material Safety Data Sheet are available on request.

WARNINGS AND PRECAUTIONS

USE BIOCIDES SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE, AND REFER TO ORGANISATIONAL POLICY AND THE SAFETY DATA SHEET.



Avoid contact with skin and eyes.



Do not breathe vapour, mist, spray, fume, dust or gas.



Make up and use with adequate ventilation.



Always use cold water to make ChloroSan solution.



DO NOT use directly on urine or vomit. Mixing this product with acid or ammonia releases chlorine gas.



Avoid prolonged contact with stainless steel, clothing or soft furnishings.



In accordance with COSHH regulations, keep tablet container tightly closed in a secure, dry, locked location, out of the reach of children.



Never remove tablet labels.



Always dispose of residual solution after 24 hours, refer to your hospital policy and SDS.



Dispose of used materials in correct waste stream and wash hands thoroughly.

Scotmas Group
Purify • Protect • Perform

Pinnaclehill Industrial Estate, Kelso TD5 8AU Scotland
(T) +44 (0)1573 226901 (E) enquiries@scotmas.com
www.scotmas.info/chlorosan



Training Video